

## BIOACTIVE COMPOUNDS DETERMINATION IN FERMENTED LIQUID DRAGON FRUIT (*HYLOCEREUS POLYRHIZUS*)

**Foong, J. H., Hon, W. M., & Ho, C. W.**

Faculty of Applied Sciences, UCSI University,  
No. 1, Jalan Menara Gading, UCSI Heights, Cheras 56000, Kuala Lumpur, Malaysia.

**ABSTRACT.** *Bioactive compounds have been intensively studied due to their beneficial effects on high cholesterol, cardiovascular disease and cancer. This study focused on the bioactive compounds (fatty acids, phytosterols, betacyanins and acetic acid) and other aspects (physical, microbiological, chemical and nutritional properties) of fermented liquid dragon fruit (*Hylocereus polyrhizus*) without pasteurization sample A (SA) and with pasteurization sample B (SB). Overall, SB exhibited significantly higher concentrations ( $p < 0.05$ ) of phytosterols, betacyanins, acetic acids, total phenolic content (TPC), total flavonoid content (TFC) and total flavanol assay (TFA) as compared to SA, except fatty acids; this suggested that heat treatment (75°C for 15 s) may have exerted a favourable effect on the concentration of these bioactive compounds. Also, SB showed greater ( $p < 0.05$ ) radical scavenging capacity in 2,2'-azino-bis-3-ethylbenzthiazoline-6-sulphonic acid (ABTS) assay than SA. No microorganisms were detected in SA or SB which was possibly due to restricting internal factor such as low pH (3.94-4.00), high ethanol concentration (7.31-8.92%) and the presence of organic acids. In conclusion, the overall results suggested both SA and SB were microbiologically safe for consumption however heat pasteurization may exert a beneficial effect on concentrations of bioactive compounds.*

**KEYWORDS.** Bioactive compounds, dragon fruit, cholesterol, pasteurization, fermentation

### INTRODUCTION

Over past decades, transitions in dietary habits and food consumption have been observed in developed and developing countries due to urbanization and accelerated economic status, and that has also led to the rise of diet-associated non-communicable diseases such as diabetes and hypertension at an alarming rate. The increased rates of these diseases have led to the rise of demand for functional foods, which are demonstrated to offer health benefits and reduce risk of chronic diseases beyond the widely accepted nutritional effects (Smith & Edward, 2010).

Fermented products are one of the popular functional food choices due to their good functional and nutritional properties (Stanton *et al.*, 2005). Development of new functional products can be expensive and challenging. One of the products introduced in recent years is the fermented dragon fruit drink. Red dragon fruit (*Hylocereus polyrhizus*) is well known in the fruit market as an excellent source of antioxidants and for its high content of nutrients, particularly potassium, as well as dietary fibre. Frequent consumption of red dragon fruit has been reported to improve overall digestive system, reduce cholesterol and prevent constipation. However, fresh dragon fruit can only last a maximum of 14 days at 10°C and 5 days at room temperature. Consequently, it is desirable to process fresh fruit into fermented liquid dragon fruit in order to avoid waste and increase its marketability (Bellec *et al.*, 2006).

In recent years, different brands of fermented dragon fruit drink have been introduced to the local market. These brands include Pitacacti Delight, Forliko, Klau, Sunberry. The consumption of these fermented dragon fruit drinks is claimed improve overall body functions (Chew, 2009). The protective health effects was strongly believed to be due to its bioactive compounds since these compounds are commonly found in plant based diets. Extensive studies have been done on cereals, legumes, nuts, oil, vegetables, fruits, tea and red wine to identify their bioactive compounds as well as their associated health benefits such as antioxidant, anti-carcinogen, cholesterol reducing as well as remitting other chronic diseases (Kris-Etherton *et al.*, 2002). However, there are only limited published studies on fermented dragon fruit drink. Therefore, the objective of this study is to

determine bioactive compounds that are present in fermented liquid red dragon fruit, with and without pasteurization treatment.

## MATERIALS AND METHODS

### *Raw materials*

Fresh red dragon fruits (RDF) were purchased from Great Sun Pitaya Farm Sdn. Bhd., Selangor, Malaysia while organic raw cane sugar was purchased from a local hypermarket.

### *Chemicals*


Chemical reagents used in this study were 4-(Dimethylamino) cinnamaldehyde (Sigma Aldrich, USA), 5 $\alpha$ -cholestane (Sigma Aldrich, Israel), absolute ethanol (John Kollin Corporation, UK), ABTS chromophore (Merck, Germany), aluminium chloride hexahydrate (Fisher Chemical, UK), betanin (TCI, Japan), bis (trimethylsilyl) trifluoroacetamide (Sigma Aldrich, USA), chloroform (Friendemann Schidmt, Australia), denatured ethanol (Bumi-Pharma, Malaysia), Folin Ciocalteu phenol reagent (Merck, Germany), glacial acetic acid (Merck, Germany), hydrochloric acid fuming 37% (Merck, Germany), methanol analytical grade (Fisher Scientific, UK), methanol HPLC grade (Merck, Germany), n-heptane (Merck, Germany), n-hexane (Merck, Germany), orthophosphoric acid (R&M Chemical, UK), peptone water (Merck, Germany), petroleum ether (Friendemann Schidmt, UK), plate count agar (BD, France), potassium hydroxide (Merck, Germany), potato dextrose agar (BD, France), purified nitrogen 99.999% (MOX, Malaysia), pyridine (Merck, Germany), sodium carbonate anhydrous (Merck, Germany), sodium chloride (Merck, Germany), sodium methylate (Merck, Germany) and sodium nitrite (Friendemann Schidmt, Australia). Milli-Q system (Millipore, USA) deionised water and ultra pure water was used throughout this study.

### *Sample pre-treatment*

Red dragon fruits (RDF) with an average weight of 0.7-0.8kg were rinsed with tap water and dried to remove dirt and residues. Subsequently, the skin of RDF was peeled and discarded while the flesh was cut into pieces, each with average thickness of 5mm.

### *Fermentation*

Stainless steel fermentation tanks with 2L volume were autoclaved while utensils were cleaned and rinsed with hot water prior to fermentation. First, 1.4kg of RDF and 280g of raw cane sugar were weighed. Then, raw cane sugar (RCS) and RDF were arranged layer by layer in fermentation tanks as shown in FIGURE 1 below. Lastly, the fermentation tanks were closed tightly and stored in a clean cabinet for 8 weeks at 25°C.



**Figure 1. Arrangement of RDF and RCS in fermentation tank.**

After 8 weeks storage, fermented samples were strained into 2 Schott Duhran bottles (1L), sample A (SA) and sample B (SB). SA was not subjected to any treatment while SB was pasteurized at 75°C for 15 sec. SA and SB were then divided into sterilized universal bottles for further analyses. Samples were stored at -20°C prior to analysis to ensure their freshness.

## Analyses

### a) Instrumental Analyses

#### i) Ethanol

The determination of ethanol content was done through automated injection method with slight modification from various studies (Wang *et al.* 2003; Jacobson, 2006). Sample was first diluted 40 times with ultra pure water and filtered into each vial using syringe filter of pore size 0.45 $\mu$ m (nylon) prior to injection into gas chromatography (GC). All of the samples were assayed in triplicate. Peak area of sample was recorded and calculated based on equation  $y = 712.38x$ ,  $R^2 = 0.9971$  developed from ethanol standard curve [0.1% -0.9% (v/v)].

The alcohol content of sample was determined using Agilent A7890 gas chromatography (Agilent Ins, US), equipped with a computer containing integrator software (ChemStation), a 30m DB-Wax capillary column (250 $\mu$ m id, film thickness: 0.25  $\mu$ m) and a flame ionization detector (FID) ( $H_2$ : 30ml/min and air: 400ml/min). The flow rate of nitrogen gas was set at 25ml/min while the temperature at injector port and detector was set at 250°C and 300°C, respectively, and split injection (97.5:1 split ratios) was used (0.1  $\mu$ L for each injection). Oven temperature was controlled with a temperature elevation program during analysis, which was initially set at 40°C for 5 min, increased to 250°C at a rate of 50°C/min and maintained for 0.8 min, with a total determination time of 10 min.

#### ii) Fatty acids (FA)

Lipid was extracted from sample according to Aldo *et al.* (2007) and Amanda *et al.* (2010) with slight modification. Sample (10mL) was added with 15mL of Folch reactive solution [chloroform/methanol; 2:1(v/v)]. The mixture solution was vortexed for 5 min, followed by the addition of 3.74mL of 0.73% (w/v) sodium chloride solution. Mixture was centrifuged at 2000rpm for 2 min to speed up the separation of organic and alcoholic-water layer. After centrifugation, the lower organic layer was transferred to another centrifuge tube by filtering through a glass funnel laid with 2g of anhydrous sodium sulphate to remove remaining water while the upper layer was discharged. The solid residue remaining was re-extracted by repeating the extraction process. The combined lower layer organic solution was transferred to a round-bottom flask (RBF) and evaporated under vacuum at 40°C until dry. Extracted lipid was stored at -20°C until further analysis.

Fatty acids in sample were converted to fatty acid methyl esters (FAME) through transesterification process before injecting into GC. Extracted lipid (50mg) was dissolved in 4.75mL of hexane in a screw-capped bottle, followed by the addition of 250 $\mu$ L of sodium methoxide. The mixture was vortexed vigorously for 10 sec and allowed to settle for 5 min. Subsequently, 3mL of upper hexane layer was transferred to a new screw capped bottle and 1mL of methanol was added. The mixture was again vortexed for 10 second settled for 5 min. Upper hexane layer was transferred into new screw capped bottle laid with 1 mm layer of anhydrous sodium sulphate and left to stand at room temperature for 30 min. Lastly, upper hexane layer was filtered into vial using nylon syringe filter (0.45 $\mu$ m) and injected into GC for analysis.

Sample (1 $\mu$ L) was injected into Perkin Elmer Instrument Autosystem XL gas chromatography equipped with a split-splitless injector and a flame ionization detector (FID). A polar fused silica capillary column SPTM-2380 of 30 m length, 0.25mm internal diameter and 0.25mm film thickness was used in FAME analysis. The flow rate of hydrogen gas and nitrogen gas was fixed at 38mL/min and 18psi respectively. The temperature of injector was set at 225°C while FID was maintained at 230°C. Initially, the column oven temperature was programmed to maintain at 140°C and then increased to 200°C at a rate of 4°C/min, followed by increment to 205°C at 0.5°C/min and 225°C at the rate of 1°C/min. Column temperature was held at 225°C for 2 min. The FAME peaks were identified by comparing the retention times of samples with standard 37 Component FAME mix (Supelco, USA). Analysis was done in triplicate and each peak area was computed and expressed as  $\mu$ g of fatty acid per mL of sample. The fatty acids concentration was quantified based on Equation 1 and Equation 2.

Equation 1:

$$\text{Response factor} = \text{FA in standard peak area} / \text{FA standard concentration}$$

Equation 2:

$$\text{FA concentration in sample} = \text{FA in sample peak area} / \text{Response factor}$$

### iii) *Phytosterols*

Determination of phytosterols involves a series of processes like saponification, extraction of unsaponifiables and silylation in order to convert phytosterols into volatile derivatives that can be detected by GC (Jill 2011). The extracted lipid (250mg) was weighed and added to 5mL of 6% ethanolic KOH and 100 $\mu$ L of 5 $\alpha$ -cholestane in round-bottom flask (RBF). Three anti-bumping chips were added to ensure smooth boiling. Mixture was refluxed for 1 hr at low heat to avoid overheating in the saponification process. The saponified mixture was transferred to a centrifuge tube, followed by the addition of 10mL of ultra pure water and 10mL of petroleum ether (PE). The mixture was centrifuged at 3000rpm for 15 min. After centrifugation, approximately 10mL of upper PE layer was transferred to a clean centrifuge tube. The remaining mixture was re-extracted by repeating the addition of water, PE and centrifugation. Once 30mL of PE layer were collected, 10mL of ethanolic-water solution (1:1, v/v) was added into the PE solution. The mixture was vortexed for 10sec and centrifuged at 4500rpm for 5 min. The upper PE layer was again transferred to a new centrifuge tube. For further purification, the new PE layer was purified by repeating the addition of ethanolic-water solution and centrifugation process twice. After the final centrifugation, the PE layer was transferred into a new boiling tube and 0.5g of anhydrous sodium sulphate was added. The entire tube was wrapped with aluminium foil and Parafilm before storing at room temperature in a cabinet overnight. The PE layer was evaporated under vacuum at 25°C until almost dry. The residual solvent was purged to dryness at 55°C with 99.9% nitrogen gas. Subsequently, 60 $\mu$ L of Bis (trimethylsilyl) trifluoroacetamide (BSTFA) and 240 $\mu$ L of pyridine were added into RBF and vortexed for 10sec. The mixture was transferred into a screw capped bottle and heated at 70°C for 45 min. It was allowed to cool down before the addition of 1.7mL of heptane. Lastly, the mixture was vortexed for 10sec and filtered into a GC vial using a nylon syringe filter (0.45 $\mu$ m).

Phytosterols was determined using Agilent A7890 gas chromatography (Agilent Ins, US) equipped with 30m DB-Wax capillary column (320 $\mu$ m id, film thickness: 0.25  $\mu$ m) and a flame ionization detector (FID) (H<sub>2</sub>: 30ml/min and N<sub>2</sub>: 30ml/min). The temperature at injector port and detector was set at 250°C and 325°C, respectively, and splitless injection was used, in which injection volume was set at 1 $\mu$ L. The initial column oven temperature was programmed at 250°C and held for 1 min, and then it was increased to 300°C at a rate of 2°C/min. Phytosterol peaks were identified by comparison with the retention time of 3 main phytosterols, which are campesterol, stigmasterol and  $\beta$ -sitosterol. The concentration of campesterol ( $y = 110824x - 363.04$ ;  $R^2 = 0.99$ ), stigmasterol ( $y = 39922x - 6.83$ ;  $R^2 = 0.99$ ) and  $\beta$ -sitosterol ( $y = 51150x - 98.03$ ;  $R^2 = 0.99$ ) were quantified based on equation developed from standard curve. Its concentration was expressed as  $\mu$ g/mL of sample.

### iv) *Organic acids and betacyanins*

Acetic acid standard solution with concentration ranging 50ppm, 100ppm, 200ppm, 400ppm and 800ppm was prepared from glacial acetic acid (Merck, Germany) while betanin standard which consists of betanin and isobetanin (500ppm, 1000ppm, 2000ppm, 4000ppm and 8000ppm) was prepared from betanin (TCI, Japan). The sample was first diluted 10 times with ultra pure water before filtering through nylon syringe filter (0.45 $\mu$ m) into sample vial.

An Agilent 1200 series HPLC system fitted with G1322A degasser, C1328B injector, G1329A column oven and G1315D diode array detector (DAD) was employed. The column used throughout this study was a reversed phase Zorbax Eclipse Plus C18 column 3.5 Micron (2.1 x 150mm). Sample injection volume was set at 10 $\mu$ L while the elution was performed at a solvent flow rate of 0.2mL/min. Temperature of column was maintained at 40°C and detection under UV spectra was carried out at 210nm (organic acids), 280nm (catechins) as well as 534nm (betacyanins) and indicated by peaks plotted in chromatograms.

The mobile phase was composed of 0.1% of ortho-phosphoric acid in ultra pure water (v/v) (Eluent A) and 0.1% of ortho-phosphoric acid in methanol (v/v) (Eluent B). The ortho-phosphoric acid was added to the solution before filtering through 0.45 $\mu$ m nylon membrane filter disc (Millipore, England) and degassed. Table 1 shows the mobile phase composition for a binary gradient condition using dual pumping system.

**Table 1. Mobile phase composition for binary gradient condition.**

Elution time (min)	Eluent A (%)	Eluent B (%)
10	98	2
58	50	50
63	50	50
73	0	100
83	0	100
93	98	2

Peaks were identified by comparing the retention times of SA and SB with retention times for acetic acid and betanin standards. The amount of acetic acid and betacyanins was determined from the peak areas at 210nm and 534nm respectively based on the equation developed from acetic acids (50 - 800ppm)  $y = 1.5966x$ ,  $R^2=0.9996$ , betanin (500 - 8000ppm)  $y = 0.154x$ ,  $R^2 = 0.9895$  and isobetanin standard curve (500 - 8000ppm)  $y = 0.1721x$ ,  $R^2 = 0.9878$ . Each sample was analysed in triplicate.

**b) Chemical analyses**

**i) Total phenolic content (TPC)**

Determination of total phenolic content was done according to Thoo *et al.* (2010) with slight modifications. Sample (1mL) of 10-fold dilution was mixed with 1mL of Folin-Ciocalteu reagent (10x dilution). After 3 min, 800 $\mu$ L of 7.5% (w/v) anhydrous sodium carbonate was added to the mixture and vortexed well. Subsequently, the mixture was incubated in the dark at room temperature for 2 hr followed by absorbance measurement at 765nm against blank using UV-vis spectrophotometer (UviLine9400, France). Results were expressed as mg of gallic acid equivalent (GAE) per 100mL of sample based on the equation  $y = 0.0396x$ ,  $R^2 = 0.9975$  using gallic acid as standard.

**ii) Total flavonoid content (TFC)**

Determination of total flavonoid content (TFC) was performed according to procedures described by Ozsoy *et al.* (2007) and Thoo *et al.* (2010) with slight modifications. A sample (0.25mL) was added to 1.25mL deionized water. At time zero, 75 $\mu$ L of 5% sodium nitrite was added into the mixture, followed by 150 $\mu$ L of 10% aluminium chloride hexahydrate 5 min later. In the next 5 min, 0.5mL of 1M sodium hydroxide and 275 $\mu$ L deionized water were added and mixed well. Immediately, absorbance of reaction mixture was measured at 510nm against blank and total flavonoid content was expressed as catechin equivalent (CE) per 100mL of sample based on the equation  $y = 0.0033x$ ,  $R^2 = 0.9991$  using catechin (50-800 $\mu$ g/mL) as standard.

**iii) Total flavanol assay (TFA)**

Total flavanol assay was determined using p-dimethylaminocinnamaldehyde (DMACA) method as proposed in various studies (Ferreira *et al.*, 2004 ; Silva *et al.*, 2007; Taylor & Giusti, 2010). The sample (160 $\mu$ L) was mixed with 2.4mL of DMACA solution [2% in 2N HCL in methanol (1:1, v/v)]. The mixture was vortexed and allowed to react at room temperature for 12 - 15 min. Absorbance of reaction mixture at 640nm was measured against a blank (DMACA solution). The concentration of TFA was calculated based on the equation  $y = 0.0022x + 0.039$ ,  $R^2 > 0.99$  using catechin as standard. Results were expressed as mg catechin equivalent (CE) per 100mL of sample.

**iv) ABTS radical scavenging capacity**

ABTS radical solution was prepared gently by mixing 10mL of 7mM ABTS solution with 10mL of 2.45mM potassium persulphate solution. The mixture was stored in a cabinet at room temperature for 12 - 16 hr before use. The ABTS radical solution was adjusted with ethanol to an absorbance  $0.7 \pm 0.02$  at 734nm before use. The sample (25 $\mu$ L) was added to 975 $\mu$ L of ABTS radical solution and allowed to react for 6 min. The absorbance of the sample was measured at 734nm against blank. The percentage of ABTS radical scavenging capacity was calculated as  $[1 - (A_e / A_c)] \times 100\%$  ( $A_e = A_{734}$  in

the presence of sample;  $A_c = A_{734}$  of negative control solution). The ABTS was expressed as The ABTS was then expressed as micromoles of trolox equivalents antioxidant capacity (TEAC) using an equation obtained from standard curve of trolox (0.1–0.8 mM)  $y = 120.1142x$ ,  $R^2 = 0.9984$  (Thoo *et al.*, 2010).

**c) Physical analyses**

**i) pH**

The pH of SA and SB were measured using pH meter (Mettler Toledo, Switzerland). Calibration was done using buffer solution of pH7 and pH4 prior to measurement. pH of sample was measured by immersing the probe into sample. Readings were recorded once there was no fluctuation shown.

**ii) Total soluble solids (TSS)**

Hand-held refractometer with detection Brix° range of 0 – 32 Brix° (Fisher Scientific, UK) was used to determine TSS of samples. Samples were equilibrated to room temperature before use. Refractometer was first calibrated with distilled water and a reading of 0 Brix° obtained. A few drops of sample were placed on the prism surface with daylight plate closed gently. Refractometer was held in direction of natural light and reading was recorded (Vasquez & Mueller, 2009).

**d) Microbiological analyses**

Microbiological analyses were aimed at evaluating the safety of food based on the presence of micro-organisms. The microbiological analyses performed were total plate counts (TPCs) and yeast and moulds (YM). According to Centre for Food Safety (2007) and NSW Food Authority (2009), the safety limit of TPC for ready-to-eat foods is  $10^4$  to  $10^5$  CFU/g generally. Triplicates were done for each microbiological analysis for SA and SB.

**i) Total plate counts (TPCs)**

Plate count agar was prepared from Difco™ Plate Count Agar powder (BD, France). A serial dilution of  $10^{-1}$ ,  $10^{-2}$  and  $10^{-3}$  were prepared. TPCs assay was carried out using spread plate method and incubated at 37°C in incubator for  $48 \pm 2$  hours. Any light amber, slightly opalescent colonies formed were recorded (Christison *et al.*, 2007; Dolci *et al.*, 2008).

**ii) Yeast and moulds (YM)**

Potato dextrose agar (PDA) was prepared from Difco™ Potato Dextrose Agar powder (BD, France) for detection of YM in SA and SB. 1mL of 10% tartaric acid was added to into every 100mL of media solution before media solidified. Dilution factor of 0,  $10^{-1}$  and  $10^{-2}$  were prepared. 0.1mL of sample from each dilution was aseptically transferred to Petri dish and spread evenly. Petri dish was incubated at 25°C in cabinet for 5 days. Any light amber, clear and slightly opalescent colonies were recorded (Ukuku & Sapers, 2006).

**Statistical analysis**

All experiment results were analysed using Minitab software (Minitab version 15.1.1.0). All numerical data were expressed as mean  $\pm$  standard deviations. The repeatability of these experiment results was indicated with relative standard deviation expressed in percentage (%RSD) as shown in Equation 3. Significant differences ( $p < 0.05$ ) between the means were determined using one-way analysis of variance (ANOVA) with Tukey's test.

Equation 3:

$$\%RSD = (\text{Standard deviation}/\text{Mean}) \times 100$$

## RESULTS AND DISCUSSION

*Repeatability***Table 2. RSD (%) of fermented dragon fruit liquid sample A (SA) and sample B (SB).**

<b>Analyses</b>	<b>SA</b>	<b>SB</b>
<b><u>Physical</u></b>		
pH	0.15	0.14
Total soluble solids	0.50	0.51
<b><u>Microbiological</u></b>		
Total plate count	N.A	N.A
Yeast and moulds	N.A	N.A
<b><u>Gas chromatography (GC)</u></b>		
Ethanol	1.09	0.47
<b><u>Fatty acids</u></b>		
Stearic acid (C16:0)	6.36	1.66
Oleic acid (C18:1n9c)	0.28	7.25
<b><u>Phytosterols</u></b>		
Campesterol	0.17	0.04
Stigmasterol	N.A	6.89
B-sitosterol	N.A	0.46
<b><u>High performance liquid chromatography (HPLC)</u></b>		
Acetic acid	4.38	6.00
Betanin	2.11	4.27
Isobetanin	1.44	4.25
<b><u>Chemical analyses</u></b>		
Total phenolic content (TPC)	0.04	0.89
Total flavonoid content (TFC)	2.25	1.80
Total flavanol assay (TFA)	0.86	1.35
ABTS radical scavenging activity	0.84	0.50

N.A Not applicable

RSD Relative standard deviation

The precision of physical, microbiological, GC, HPLC and chemical analyses were evaluated in terms of relative standard deviation (RSD) as shown in TABLE 2. Overall, the RSD ranged from 0.00 to 7.25. These low values of RSD below 10% indicated the small variability and high precision of all physical, microbiological, instrumental and chemical analyses (Bievre & Gunzler, 2005; California Department of Toxic Substance, 2006).

As compared to other analyses, GC analyses have a slightly higher range of RSD values from 0.04 to 7.25. One of the possible factors was manual injection of sample into GC in fatty acid analyses because injection accuracy, speed of injection and syringe handling technique are crucial in accurate quantisation of fatty acid methyl esters (FAME). It is possible that some loss of the high boiling components may have occurred during initial injection entry and also after the removal of needle due to vaporization and fractionation of sample from needle (Barwick, 1999). Apart from that, pre-treatment of sample prior to injection into GC may contribute to larger RSD as these are subjective to personal handling techniques and consistency.

**Determination of Fatty Acids and Bioactive Compounds****Table 3. Fatty acids and bioactive compounds in sample A (SA) and sample B (SB).**

<b>Analyses</b>	<b>SA<sup>x</sup></b>	<b>SB<sup>x</sup></b>
<b><i>GC analysis</i></b>		
<b><i>Fatty acids</i></b>		
Stearic acid (C18:0) (mg/L)	2.00±0.13 <sup>b</sup>	1.52±0.03 <sup>a</sup>
Oleic acid (C18:1n9c) (mg/L)	5.10±0.01 <sup>a</sup>	3.52±0.26 <sup>b</sup>
<b><i>Phytosterols</i></b>		
Campesterol (mg/L)	3.38±0.01 <sup>b</sup>	3.64±0.00 <sup>a</sup>
Stigmasterol (mg/L)	N.D	0.82±0.06 <sup>a</sup>
B-sitosterol (mg/L)	N.D	2.83±0.01 <sup>a</sup>
<b><i>HPLC analysis</i></b>		
Acetic acid (g/L)	5.15±0.23 <sup>b</sup>	6.52±0.39 <sup>a</sup>
Betanin (g/L)	13.24±0.28 <sup>a</sup>	14.23±0.61 <sup>a</sup>
Isobetanin (g/L)	7.31±0.11 <sup>b</sup>	12.73±0.54 <sup>a</sup>
<b><i>Chemical analysis</i></b>		
Total phenolic content (TPC) (mg/100mL GAE)	371.63±0.15 <sup>b</sup>	396.12±3.61 <sup>a</sup>
Total flavonoid content (TFC) (mg/100mL CE)	119.70±2.69 <sup>b</sup>	140.51±2.52 <sup>a</sup>
Total flavanol assay (TFA) (mg/100mL CE)	81.21±0.69 <sup>b</sup>	100.91±1.36 <sup>a</sup>

<sup>x</sup> Each value in table represents the mean value ± standard deviation

<sup>a-b</sup> Mean value with different superscript in each column differs significantly ( $p < 0.05$ )

N.D Not detected

GAE Gallic acid equivalent

CE Catechin equivalent

Bioactive compounds refer to essential and non-essential compounds that occur in nature, are part of the food chain, and can be shown to have an effect on human health (Bielsalski *et al.*, 2009). They typically occur in small quantities and present as natural constituents in food that provide health benefits beyond the basic nutritional value of the product. Bioactive compounds have been intensively studied lately due to their beneficial effect on cholesterol, cardiovascular diseases and cancer shown in many studies (Kris-Etherton *et al.*, 2002). These bioactive compounds are usually divided into various groups based on their chemical structure and functions like phenolic compounds and its subcategory, flavonoids, antioxidants pigments like betalains and its subgroup betacyanins (Henriette, 2009) as well as phytosterols (Saura & Goni, 2009).

Apart from bioactive compounds, two main fatty acids, namely stearic acid and oleic acid, were emphasized in SA and SB due to their beneficial effect on cholesterol level. Stearic acid was reported to have “neutral” effect on LDL and cholesterol level due to its rapid transformation into oleic acid (Valentin *et al.*, 2005) while Craig *et al.* (2000) suggested that stearic acid reduced plasma cholesterol level by reducing cholesterol absorption and increased excretion of endogenous cholesterol in hamster study. Oleic acid is well-studied for its effectiveness in reducing cholesterol level and decreases the risk of cardiovascular disease as reported by Stoutjesdijk *et al.* (2000), Kalyana *et al.* (2003) and Ribeiro *et al.* (2009). The results obtained were in accordance with Khor (2011) who also found that stearic acid and oleic acid are the main fatty acids in red dragon fruits (RDF) seeds. It was suggested that fatty acids and phytosterols found in RDF seed penetrated out from seeds into fermented liquid during fermentation process. This is supported by Fadahunsi & Sunni (2010) as well as Afolabi *et al.* (2011) who mentioned that fermentation improved the fatty acids composition and oil yield from seeds. SA has significantly higher concentration ( $p < 0.05$ ) of stearic acid and oleic acid than SB, which were 2.00mg/L and 5.10mg/L respectively. This suggested

that heat pasteurization on SB caused the reduction in stearic acid and oleic acid concentration, which is in accordance with Agata *et al.* (2005) and Somboonsilp *et al.* (2011) who also reported the decrease of stearic acids in their studies of heat pasteurization on milk and juice respectively.

Phytosterols are natural components that are largely derived from vegetable oils, plants and cereals. Among various plant sterol structures that have been discovered, the phytosterols found in the highest abundance in most plants are sitosterol, campesterol and stigmasterol (Pascal & Segal, 2006). According to Isabelle *et al.* (2009), recommended daily intake of 2g of phytosterols reduced LDL-cholesterol concentration by 10% by inhibiting cholesterol absorption by 30-40%. Therefore, phytosterols fortified food products were given much interest. From Table 3, SB showed a significantly higher concentration ( $p < 0.05$ ) of phytosterols (campesterol, stigmasterol and  $\beta$ -sitosterol) in contrast to fatty acids as compared to SA. Stigmasterol and  $\beta$ -sitosterol were not detected in SA probably due to its low concentration that was beyond the detection limit. One of the possible reasons of lower phytosterol concentration in SA was oxidation. According to Soupas (2006), phytosterols are prone to oxidation due to the double bond at the 5-position and the percentage of oxidation was dependent on the lipid matrix, fatty acids composition and sterol content. For instance, oxidation of sitosterol was noted to increase with increasing unsaturation of lipid matrix. This probably explained the lower concentration of phytosterols in SA due to its higher content of unsaturated fatty acids. The absence of stigmasterol and  $\beta$ -sitosterol was also supported by Carreno *et al.* (2008) who pointed out that stigmasterol, followed by cholesterol and sitosterol has the greatest susceptibility to oxidation in oil-in-water emulsions due to their differences in their surface activity. Also, it was suggested that the pasteurization temperature and time were too short to cause the significant loss of phytosterols in SB. This was supported by Carren *et al.* (2008) who reported that milk subjected to Schaal oven condition at 65°C for 24 hours only slightly reduced phytosterol content by 4%. In addition, Linda (2002) also stated that chemical stability of heat pasteurized phytosterol-esters containing milk and yogurt were found to be similar to standard product.

Organic acid is one of the common byproducts formed during fermentation. The content of organic acid not only influences the flavours but also the stability, acceptability, nutrition and keeping quality of a product (Shui & Leong, 2002). Acetic acid was emphasized in this study due to its beneficial effect on lowering cholesterol level as reported by Fushimi *et al.* (2006) and Dash (2009) on cholesterol-fed rats. As analysed by HPLC, SB showed a significantly higher acetic acid concentration compared to SA. The possible explanation was the exposure of SB to heat pasteurization increased the rate of acetate esters hydrolysis. This was supported by Pascal *et al.* (2006) who stated that ethyl acetate is one of the most prevalent esters found in fermented wine. The breakdown of esters thus increased the concentration of acetic acid in liquid (Jackson, 2008). The result was further supported by the higher ethanol concentration of SB as shown in Table 5.

Besides that, SB was found to exhibit significant higher concentration ( $p < 0.05$ ) of isobetanin content as compared to SA. This showed that the effect of pasteurization (75°C for 15s) did not decrease the concentration of betanin and isobetanin in SB. Instead, it was likely that the temperature was not too high and the time was too short to cause significant degradation of betacyanins. This is supported by (Henriette, 2009) who reported that loss of betacyanin in dragon fruit juice acidified to pH 4 is less than 10% during pasteurization at 80°C for 5 min. Also, the low pH of SB at pH 4.0 and the presence of vitamin C naturally present in dragon fruit were reported by Henriette 2009 to improve the stability of betacyanin pigments. Therefore, the difference in betanin and isobetanin concentration of SA and SB was more likely to be caused by some other extrinsic factors like exposure to oxygen and light during sample treatment. Light and oxygen have been identified as the major factor of colour deterioration instead of heat; each caused betalain degradation by 15.6% and 14.6% as reported by Woo *et al.* (2011). In addition, different structures of betalains also contributed to its stability. For instance, glycosylated structures are more stable than a glycons, probably because of the higher oxidation–reduction potentials of the former (Henriette, 2009). Since betacyanins are most often glycosylated, thus it is likely to be stable when subjected to heat pasteurization for a short time (Mortensen, 2006).

Betacyanin was given much interest lately due to its stability as a natural food colorant as well as its excellent antioxidant properties (Rebecca *et al.*, 2010; Woo *et al.*, 2011). In addition, Tsai *et al.* (2011) also reported its potent beneficial effect of lowering cholesterol. There are at least seven identified betacyanins in *Hylocereus polyrhizus*, in which betanin, isobetanin, betanidin, isobetanidin

being the major red-violet pigment and others such as phylloactin, isophylloactin, and bougainvillein-R-I, where all have identical absorption spectra that contribute to the colour of pulp (Henriette, 2009; Rebecca *et al.*, 2010). Among these, only betanin and isobetanin were quantified in this study due to the unavailability of commercial standard.

Plant source food bioactive compounds such as phenolic compounds have been recognized as having potential to reduce disease risk such as coronary heart disease, cancer and reduce cholesterol level, mainly arising from its antioxidant properties (Yang *et al.*, 2007; Kuraswamy & Satish, 2008; Zainoldin & Baba, 2012). Therefore, various spectrophotometric methods were used for the quantification of phenolic compounds in fermented RDF liquid, such as total phenolic content (TPC), total flavonoid content (TFC) and total flavanol assay (TFA). All of these assays are used to determine various structural groups phenolic compounds based on different principles (Chang *et al.*, 2002; Tay, 2011). From Table 3, SB was found to exhibit significant higher ( $p < 0.05$ ) value of TPC, TFC and TFA as compared to SA.

Total phenolic content (TPC) assay measures both free and bound phenolic compounds. The significant higher value of TPC in SB could be explained by the higher concentration of betacyanins that present. This is because betacyanins could also contribute to the total phenolic due to the presence of phenol structure in the molecule (Zainoldin & Baba, 2012). Apart from that, Anup *et al.* (2011) also found that pasteurized aonla juice at 75°C had a higher amount of gallic acid during storage. It was likely that the effect of pasteurization also contributed to the increased concentration of total phenolics compounds in SB. This was supported by Yang *et al.* (2007) who reported that pasteurization at 65°C and 75°C will not significantly decrease noni juice quality as most of the phenolics from plants present in a more stable conjugated form with glycosides or esters. In addition, the difference in TPC value between SA and SB may also be affected by the presence of other compounds like oleic acid and reducing sugars like fructose (Yu, 2008).

Red dragon fruits are rich in naturally-occurring flavonoids (Kunnika & Pranee, 2011). These flavonoids were reported to have a wide range of biological activities such as antioxidant, enzyme-inhibiting, and antibacterial effects (Lee, 2009). Total flavonoid content (TFC) was examined by aluminium chloride colorimetric method. This method does not detect all kinds of flavonoids, instead only flavones and flavonols were found to complex stably with aluminium chloride. Even though it is not suitable to obtain values of total flavonoids, it is sufficient for comparative analysis (Chang *et al.*, 2002). In contrast to Laura *et al.* (2010) and Igual *et al.* (2011) who reported that heat pasteurization generally reduced flavonoid content in juices, SB showed a significantly higher ( $p < 0.05$ ) TFC value than SA. Hence, it is suggested that the difference in TFC value was due to the variation in ripeness of red dragon fruit in SA and SB.

Total flavanol assay (TFA) is mainly used to determine flavanols such as epigallocatechin gallate (EGCG), epigallocatechin (EGC), epicatechin gallate (ECG), epicatechin (EC) and catechin (C), which are well-known for their strong antioxidant properties (Cabrera *et al.* 2006). Among these, affinity of DMACA to (-)-epicatechin was reported to be a few folds higher than other individual flavanol compounds (Treutter, 1989). As shown in TABLE 3, SB showed a significantly higher ( $p < 0.05$ ) TFA value than SA. This was in accordance with results reported by Fuleki and Ricardo (2003) who mentioned that pasteurization increased the concentration of catechins in cold-pressed juice. Also, Zimmerman & Gleichenhagen (2011) reported that higher steeping temperature of green tea infusions resulted in higher amount of catechins. Thus, it was suggested that the higher concentration of TFA in SB was due to the higher rate of catechins diffused into liquid as the pasteurization temperature at 75°C was not too high to cause significant degradation of catechins. This was supported by Ziaedini *et al.* (2010) who observed that catechin degradation occurred in the temperature range of 80-90°C.

**Antioxidant Scavenging Capacity****Table 4. Antioxidant scavenging capacity of sample A (SA) and sample B (SB).**

Chemical analyses	Unit	SA <sup>x</sup>	SB <sup>x</sup>
ABTS radical scavenging capacity	mM TEAC	0.54±0.02 <sup>b</sup>	0.59±0.01 <sup>a</sup>

<sup>x</sup> Each value in table represents the mean value ± standard deviation

<sup>a-b</sup> Mean value with different superscript in each column differs significantly ( $p < 0.05$ )

TEAC Trolox equivalent antioxidant capacity

ABTS radical-scavenging capacity assay was chosen instead of the commonly used DPPH assay due to the absorbance interruption at 517 nm by the similar purple colour of fermented RDF liquid. In ABTS radical scavenging capacity assay, ABTS was converted to its radical cation by the addition of potassium per sulphate. The radical cation is reactive towards most antioxidants which include phenolics, thiols, and vitamin C (Bhagat *et al.*, 2011). As shown in Table 4, SB exhibited a significantly higher ( $p < 0.05$ ) radical scavenging capacity than SA. This was probably due to the significantly higher concentration of betanin, isobetainin, TPC, TFC as well as TFA of SB than in SA. Even though SA showed a higher vitamin C concentration than SB, it was suggested that betacyanins exhibited stronger oxidative power than vitamin C. These were supported by Cai *et al.* (2003) who reported that betalains showed stronger antioxidative power than ascorbic acid and catechins in the plants of *Amaranthaceae* and Tenore *et al.* (2011) who found that betacyanins fractions exhibited the highest radical-scavenging capacity among flavonoids and phenolics in red pitaya fruits. Also, phenolic content was reported by Lee *et al.* (2003) to be highly correlated with antioxidant capacity from ABTS assay and Lee (2009) mentioned that phenolic and polyphenolic compounds constituted the main class of natural antioxidants present in plants that accounted for their strong antioxidant properties. Besides that, heat pasteurization may also be one of the factors that contributed to higher radical scavenging capacity as heat treatment may release bound antioxidants as free antioxidants as stated by Yang *et al.* (2007). The cause of variations in the ABTS assay may be that the ABTS working solution was not always the same age, hence the reaction of the solution with fermented RDF liquid might have been different during the determination time (Thaipong *et al.*, 2006) and the ABTS radical solution concentration is difficult to control (Cai *et al.*, 2003).

**Physico-Chemical Properties****Table 5. Physico-chemical profile of sample A (SA) and sample B (SB).**

Analyses	Unit	SA <sup>x</sup>	SB <sup>x</sup>
pH	-	3.94±0.01 <sup>b</sup>	4.00±0.01 <sup>a</sup>
Total soluble solids (TSS)	Brix <sup>o</sup>	23.1±0.12 <sup>a</sup>	22.7±0.12 <sup>b</sup>
Ethanol	%	7.31±0.08 <sup>b</sup>	8.92±0.04 <sup>a</sup>

<sup>x</sup> Each value in table represents the mean value ± standard deviation

<sup>a-b</sup> Mean value with different superscript in each column differs significantly ( $p < 0.05$ )

Physico-chemical analyses are commonly used in determining food stability because physical properties of a food affect the environment for chemical reactions and microbial growth. As shown in Table 5, it was found that SA and SB were significantly different ( $p < 0.05$ ) from the aspect of pH, TSS and ethanol content. This suggested that the pasteurization treatment of SB may be one factor that caused the significant difference ( $p < 0.05$ ) in the physical properties of fermented RDF liquid as compared to non-pasteurized SA. However the results obtained were in contrast with some of the previous findings which stated that no remarkable increase in pH was observed after different thermal treatment by Rivas *et al.* (2006) and Sampedro *et al.* (2009). Therefore, it was more likely that the difference in pH was due to the different amount of organic acids produced from two individual fermentation tanks. Even so, the pH range of SA and SB were found to be acceptable and similar to

some other fermented fruit wine like palm sap wine (pH 4.0), coconut palm sap (pH 3.7), Jambal fruit wine (pH 3.5), mango wine (pH 3.7) and banana wine (pH 3.85) as reported by George (2008).

Total soluble solids (TSS) is a measure of overall sugar content of mixture. Besides being a parameter in assessing food physical stability, it is also often associated with food sweetness and consumer acceptability (Harker *et al.*, 2002). Sampedro *et al.* (2009) reported no significant variation in Brix<sup>o</sup> for thermally treated orange juice. It is more likely that the significant difference in TSS of SA and SB was due to the difference in ripeness of fruit as higher Brix<sup>o</sup> is found in riper fruit as suggested by Rivas *et al.* (2006).

Ethanol is one of the by-products formed from fermentation particularly by heterogenic lactic acid bacteria and yeasts as stated by Asli (2010). It is believed to be the main contributor for the alcoholic flavour in SA and SB due to its presence in large quantities as well as its high reactivity of primary alcohols (Chung *et al.*, 2008; Sumbly *et al.*, 2010). Overall, SB has significant higher ethanol content than SA. The possible reason was the exposure of SB to high heat during pasteurization. According to Jackson (2008), temperature directly affects the rate of reactions particularly ester hydrolysis in wine during storage. This may explain the significantly ( $p < 0.05$ ) higher ethanol (Table 5) and acetic acid content (Table 3) in SB as accelerated ester hydrolysis resulted in more ethanol and organic acids being released into the sample (Sumbly *et al.*, 2010). These results were also supported by Patricia *et al.* (2006) who reported that acetate esters hydrolysed more quickly than other ethyl esters and these significant decreases were found in wine subjected to higher temperature at 45°C.

### Microbiological Properties

**Table 6. Microbiological profile of sample A (SA) and sample B (SB).**

Microbiological analyses	Unit	SA	SB
Total plate count	CFU/mL	N.D	N.D
Yeast and mould	CFU/mL	N.D	N.D
N.D	Not detected		
CFU	Colonies forming unit		

Microbiological analysis plays an important role in judging the quality and safety of a food product. The growth of microorganisms and presence of pathogenic microorganisms need to be considered as these will lead to food spoilage and food safety issues (Yousef & Calstrom, 2003). As shown in TABLE 6, no growth of aerobic mesophilic bacteria, yeast or moulds was observed in either SA or SB. This indicated that SA and SB fall under the satisfactory level of microbiological standards for ready-to-eat food, which is less than 10<sup>4</sup> CFU/mL of sample (Centre for Food Safety, 2007; NSW Food Authority, 2009). Therefore both SA and SB can be confirmed microbiologically safe for consumption.

The absence of microorganisms regardless of pasteurization could be due to the restricting internal factor of SA and SB such as low pH, presence of ethanol, carbon dioxide and organic acids that were formed during fermentation as suggested by Erbas *et al.* (2005). Similar results were reported in other fermented products like plant beverage and maize by Kantachote *et al.* (2008), Aloba & Offonry (2009), Namugumya & Muyanja (2009), Chelule *et al.* (2010) who also reported that the combined effect of low pH (3.5-4.0) and presence of organic acids and ethanol were found inhibitive to bacteria.

## CONCLUSION

In summary, these results confirmed that both SA and SB were microbiologically safe for consumption due to the absence of micro-organisms. The effect of pasteurization suggested possible beneficial effects on bioactive compound concentration as SB showed significantly ( $p < 0.05$ ) higher amounts of phytosterols (campesterol, stigmasterol,  $\beta$ -sitosterol), betacyanins (betanin, isobetanin), acetic acids, total phenolic content (TPC), total flavonoid content (TFC) and total flavanol assay

(TFA) as well as greater ( $p < 0.05$ ) radical scavenging capacity in ABTS assay as compared to SA. However, the pasteurization also reduced the concentration of fatty acids, vitamins and minerals in SB. Further study can be done to maximize the concentration of bioactive compounds while minimizing the loss of vulnerable vitamins and minerals.

### ACKNOWLEDGEMENT

Firstly, I would like to express my sincere appreciation to Malaysia Ministry of Higher Education for the Fundamental Research Grant Scheme (FRGS) monetary funding (FRGS/2010/2/STWN/UCSI/03/1) to carry out this project. Also, I am very thankful to my supervisor and laboratory staff, especially Mr. Kok, Mr. Lingam, Mr. Chua, Ms. Ang, Ms. Vasuki, Mr. Neo, Ms. Choi and Ms. Tan for their technical assistance their valuable guidance, constructive comments, encouragement and insight throughout this research.

### REFERENCES

- Afolabi, I. S., Marcus, G. D., Olanrewaju, T. O., & Chizea, V. 2011. Biochemical Effect of Some Food Processing Methods on the Health Promoting Properties of Under-Utilized Carica Papaya Seed. *Journal of Natural Products*, **4**: 17-24.
- Agata, R., Genowefa, B., & Pustkowiak, H. 2005. Effect of Heat Treatment on the Free Fattyacids of Ewe's Milk. *Biotechnology I Animal Husbandry*, **21 (6)**: 237-240.
- Aldo, P., Samantha, S., Gino, T., Nico, B., & Gianfranco, P., 2007. Different Level of Conjugated Linoleic Acid (CLA) in Dairy Products from Italy. *Journal of Food Composition and Analysis*, **20**: 472-479.
- Alobo, A. P., & Offonry, S. U. 2009. Characteristics of Coloured Wine Produced from Roselle (*Hibiscus sabdariffa*) Calyx Extract. *Journal of the Institute of Brewing*, **115 (2)**: 91-94.
- Amanda, T., Beatriz, C., Salvador, M., José, L. T., & Soledad, P. 2010. Evaluation of Different Fat Extraction and Derivatization Strategies for the Determination of the Fatty Acid Profile of Ice Creams by Gas Chromatography. *Food Innovation*, pg 1-4.
- Amran, A., Zakaria, Z., Othman, F., Das, S., Raj, S., & Nordin, M. 2010. Aqueous Extract of *Piper sarmentosum* Decreases Atherosclerotic Lesions in High Cholesterolemic Experimental Rabbits. *Lipid in Health and Disease*, **44 (9)**: 1-6.
- Anup, K. B., Tandon, D. K., Dikshit, A., & Kumar, S. 2011. Effect of Pasteurization Temperature on Quality of Aonla Juice during Storage. *Journal of Food Science and Technology*, **48 (3)**: 269-273.
- Asli, M. A. 2010. A Study on some Efficient Parameters in Batch Fermentation of Ethanol using *Saccharomyces cerevesiae* SC1 Extracted from Fermented Siahe Sardasht Pomace. *African Journal of Biotechnology*, **9 (20)**: 2906-2912.
- Augustin, S., & Gary, W. 2000. Dietary Intake and Bioavailability of Polyphenols. *Journal of Nutrition*, **130**: 20735-20855.
- Australian Government. 2006. *Vitamins and Minerals* [Online]. Available from: <http://www.health.gov.au/internet/healthyactive/publishing.nsf/content/vitamins-minerals> [Accessed 22 December 2011].
- Barwick, V. J. 1999. Sources of Uncertainty in Gas Chromatography and High-Performance Liquid Chromatography. *Journal of Chromatography A*, **849**: 13-33.
- Bellec, F., Vaillant, F., Imbert, W. H., Campbell, M. K., & Farrell, S. O. 2009. *Introduction to General, Organic and Biochemistry*, 9<sup>th</sup> ed. US: Cengage Learning, p.485.
- Bhagat, J., Lobo, R., Parmar, V., & Ballal, M. 2011. *In vitro* Free Radical Scavenging Potential of Indian Habitant *Anisochilus carnosus* (L..F.) Wall. *Chinese Journal of Natural Medicines*, **9 (6)**: 456-460.
- Biesalski, H., Dragsted, L. O., Ibrahim, E., Grossklaus, R., Muller, M., Schrenk, D., Walter, P., & Weber, P. 2009. Bioactive Compounds : Definition and Assessment of Activity. *Journal of Nutrition*, **25 (11)**: 1202-1205.

- Bievre, P. D., & Gunzler, H. 2005. *Validation in Chemical Measurement*. New York: Springer, p. 4.
- Cabrera, C., Artachon, R., & Gimenez, R. 2006. Beneficial Effects of Green Tea: A Review. *Journal of the American College of Nutrition*, **25** (2): 79-99.
- Cai, Y. Z., Sun, M., & Corke, H. 2003. Antioxidant Activity of Betalains from Plants of the Amaranthaceae. *Journal of Food Agriculture Food Chemistry*, **51**: 2288-2294.
- California Department of Toxic Substance. 2006. *Interpretation of Analytical Data* [Online]. Available from: [http://www.dtsc.ca.gov/AssessingRisk/ECL/upload/6\\_06\\_UM.pdf](http://www.dtsc.ca.gov/AssessingRisk/ECL/upload/6_06_UM.pdf) [Accessed 12 December 2011].
- Carren, M. M., Diana, A., & Iciar, A. 2008. Stability of Sterols in Phytosterol-Enriched Milk under Different Heating Conditions. *Journal of Agriculture and Food Chemistry*, **56** (21): 9997-10002.
- Centre for Food Safety. 2007. Microbiological Guidelines for Ready-to-eat Food [Online]. Available from: <http://blpd.dss.go.th/micro/ready.pdf> [Accessed 21 November 2011].
- Chang, C. C., Yang, M. H., Wen, H. M., & Chern, J. C. 2002. Estimation of Total Flavonoid Content in Propolis by Two Complimentary Colorimetric Methods. *Journal of Food and Drug Analysis*, **10** (3): 178-182.
- Chelule, P. K., Mokoena, M. P., & Gqaleni, N. 2010. Advantages of Traditional Lactic Acid Bacteria Fermentation of Food in Africa. *Current Research, Technology and Education Topics in Applied Microbiology and Microbial Biotechnology*, 1160-1167.
- Chew, M. 2009. Fruit "Tonic" from the Dragon Fruit. *The Star Publication*. April 21, 2009.
- Chien, Y. L., Wu, L. Y., Lee, T. C., & Lucy, S. H. 2010. Cholesterol-lowering Effect of Phytosterol-Containing Lactic-Fermented Milk Powder in Hamsters. *Food Chemistry*, **119**: 1121-1126.
- Christison, C. A., Lindsay, D., & Holy, A. 2007. Microbiological Survey of Ready-to-eat Foods and Associated Preparation Surfaces in Retail Delicatessens, Johannesburg, South Africa. *Food Control*, **19**: 727-733.
- Chung, H. J., Son, J. H., Park, E. Y., Kim, E. J., & Lim, S. T. 2008. Effect of Vibration and Storage on some Physico-Chemical Properties of a Commercial Red Wine. *Journal of Food Composition and Analysis*, **21**: 655-659.
- Craig, L. S., Russell, L. C., Cindy, L. S., & Timothy, P. C. 2000. Dietary Stearic Acid Reduces Cholesterol Absorption and Increases Endogenous Cholesterol Excretion in Hamsters Fed Cereal-Based Diets. *Journal of Nutrition*, **130**: 1232-1238.
- Dash, A. T. 2009. *Secrets to Lowering Cholesterol with Nutrition and Natural Supplements, Safely*. 1<sup>st</sup> ed. US: Author House, p. 90.
- Dillard, A., Matthan, N., & Lichtenstein, A. H. 2010. Use of Hamsters as a Model to Study Diet-Induced Atherosclerosis. *Nutrition and Metabolism*, **89** (7): 1-12.
- Ditscheid, B., Keller, S., & Jahreis, G. 2005. Cholesterol Metabolism is affected by Calcium Phosphate Supplementation in Humans. *The American Society for Nutritional Science*, **135**: 1678-1682.
- Dolci, P., Alessandria, V., Zeppa, G., Rantsiou, K., & Cocolin, L. 2008. Microbiological Characterization of Artisanal Raschera PDO Cheese: Analysis of its Indigenous Lactic Acid Bacteria. *Food Microbiology*, **25**: 392-399.
- DSM Nutritional Products. 2010. *Fortification Basics* [Online]. Available from: [http://www.dsm.com/en\\_US/downloads/dnp/51610\\_fort\\_basics\\_stability.pdf](http://www.dsm.com/en_US/downloads/dnp/51610_fort_basics_stability.pdf) [Accessed 22 December 2011].
- Erbas, M., Certel, M., & Uslu, M. K. 2005. Microbiological and Chemical Properties of during Fermentation and Storage as Wet-sensorial Properties of Tarhana Soup. *Learning with Technology, Foods Science and Technology*, **38**: 409-416.
- European Union Legislation. 2010. *Maximum Level for Certain Contaminants* [Online]. Available from: [http://europa.eu/legislation\\_summaries/food\\_safety/contamination\\_environmental\\_factors/l21290\\_en.htm](http://europa.eu/legislation_summaries/food_safety/contamination_environmental_factors/l21290_en.htm) [Accessed 13 December 2011].
- Fadahunsi, I. F., & Sanni, A. I. 2010. Chemical and Biochemical Changes in Bambara Nut (*Voandzeia subterranean*) during Fermentation to 'Tempeh'. *Electronic Journal of Environmental, Agriculture and Food Chemistry*, **9** (2): 275-283.

- Ferial, S. A., Shahinaz, A. H., Abeer, M. N. H., & Ibrahim, M. A. 2011. Stability of Different Fruit Juices Mixed with Black Carrot Juice during Storage. *International Journal of Academic Research*, **3** (6): 293-301.
- Ferreira, D., Slade, D., & Maraise, J. P. J. 2006. Flavans And Proanthocyanidins. In. OM Andersen and KR Markham. Flavanoids Chemistry, *Biochemistry And Applications*. New York: Taylor and Francis.
- Food and Agricultural Organization (FAO). 2008. *Fats and Fatty Acids in Human Nutrition*. [Online]. Available from: <http://www.fao.org/docrep/013/i1953e/i1953e00.pdf> [Accessed 22 December 2011].
- Fuleki, T., & Ricardo, J. M. 2003. Effects of Cultivar and Processing Method on the Contents of Catechins and Procyanidins in Grape Juice. *Journal of Agriculture Food Chemistry*, **51** (3): 640-646.
- Fushimi, T., Suruga, K., Oshima, Y., Fukihar, M., Tsukamoto, Y., & Goda, T. 2006. Dietary Acetic Acid Reduces Serum Cholesterol and Triacylglycerols in Rats Fed a Cholesterol-Rich Diet. *Journal of Nutrition*, **95** (5): 916-924.
- George, W. B. 2008. *The Effect of Enzymatic Processing in Banana Beer and Wine*. Thesis. South Africa: Stellenbosch University.
- Harker, F. R., Marsh, K. B., Young, H., Murray, S. H., Gunson, F. A., & Walker, S. B. 2002. Sensory Interpretation of Instrumental Measurements 2: Sweet and Acid Taste of Apple Fruit. *Postharvest Biology and Technology*, **24**: 241-250.
- Henriette, M. C. A. 2009. Betalains: Properties, Sources, Applications, and Stability – A Review. *International Journal of Food Science and Technology*, **44**: 2365-2376.
- Higdon, J. 2005. *Flavonoids* [Online]. Linus Pauling Institute. Available from: <http://lpi.oregonstate.edu/infocenter/phytochemicals/flavonoids/> [Accessed 21 December 2011].
- Igual, M., Martinez, E. G., Camacho, M. M., & Navarrete, N. M. 2011. Changes in Flavonoid Content of Grapefruit Juice Caused by Thermal Treatment and Storage. *Innovative Food Science and Emerging Technologies*, **12** (2): 153-162.
- Isabelle, D., Rouyane, T. R., Henk, C. M., Guus, S. M., Linsie, M., Peter, L. Z., Johanna, M. G., & Elka, A. T. 2009. Continuous Dose-Response Relationship of the LDL-Cholesterol Lowering Effect of Phytosterol Intake. *Journal of Nutrition*, **139**: 271-284.
- Jackson, R. S. 2008. *Wine Science: Principles and Applications*. Third edition. US: Elsevier Academic Press, p. 280-312, 434, 504, 751.
- Jacobson, J. L. 2006. *Introduction to Wine Laboratory Practices And Procedures*. First edition. Germany: Birkhäuser, p. 140.
- Jill, W. M. 2011. *Gas Chromatographic Analysis of Plant Sterols* [Online]. American Oil Chemists' Society. Available from: <http://lipidlibrary.aocs.org/topics/phytosterols/index.htm> [Accessed 30 September 2011].
- Kalyana, S., Ravigadevi, S., & Tan, Y. A. 2003. Palm Fruit Chemistry and Nutrition. *Asia Pacific Journal of Clinical Nutrition*, **12** (3): 355-362.
- Kantachote, D., Charernjiratrakul, W., & Umsakul, K. 2008. Antibacterial Activities of Fermented Plant Beverages Collected in Southern Thailand. *Journal of Biological Sciences*, **8**: 1280-1288.
- Khalili, R., Norhayati, A. H., Rokijaj, M. Y., Asmah, R., Muskinah, M., & Manaf, A. 2009. Hypocholesterolemic Effect of Red Pitaya (*Hylocereus Sp.*) on Hypercholesterolemia Induced Rats. *International Food Research Journal*, **16**: 431-440.
- Khor, Y. P. 2011. Effect of Storage Duration on Fatty Acids and Phytosterols Changes in Red Dragon Fruit (*Hylocereus Polyrhizus*) Seeds at 4°C. Thesis. Malaysia: UCSI University.
- Kim, E. S., Liang, Y. R., Jin, J., Sun, Q. F., Lu, J. L., & Du, Y. Y. 2007. Impact of Heating on Chemical Compositions of Green Tea Liquor. *Food Chemistry*, **103**: 1263-1267.
- Ko, Y. J., Wu, Y. W., & Lin, W. C. 2004. Hypolipidemic Effect of *Glycine Tomentella* Root Extract in Hamsters. *The American Journal of Chinese Medicine*, **32** (1): 57-63.
- Kris-Etherton, P. M., Hecker, K. D., Bonanome, A., Coval, S. M., Binkoski, A. E., Hilpert, K. F., Griel, A. E., & Etherton, T. D. 2002. Bioactive Compounds in Foods: Their Role in the Prevention of Cardiovascular Disease and Cancer. *American Journal of Medicine*, **113**: 71-88.

- Kumaraswamy, M. V., & Satish, S. 2008. Antioxidant and Anti-Lipoxygenase Activity of *Thespesia Lampas* Dalz & Gibs. *Advances in Biological Research*, **2** (4): 56-59.
- Kunnika, S., & Pranee, A. 2011. Influence of Enzyme Treatment on Bioactive Compounds and Colour Stability of Betacyanin in Flesh and Peel of Red Dragon Fruit *Hylocereus polyrhizus* (Weber) Britton and Rose. *International Food Research Journal*, **18** (4): 1437-1448.
- Laura, A. R., Emilio, A. P., & Gustavo, A. G. 2010. *Fruit and Vegetable Phytochemicals: Chemistry, Nutritional Value and Stability*. First edition. US: John Wiley and Sons, p. 148.
- Lee, K. W., Kim, Y. J., Lee, H. J., & Lee, C. Y. 2003. Cocoa has more Phenolic Phytochemicals and a Higher Antioxidant Capacity than Teas and Red Wine. *Journal of Agriculture Food Chemistry*, **51**: 7292-7295.
- Lee, Q. H. 2009. Total Phenolic Content and Total Flavonoid of *H. Polyrhizus* Waste Extract by using Ultrasonic Solvents Extraction. Thesis: University Malaysia Pahang.
- Lim, T. H. 2004. Betacyanins From *Hylocereus Undatus* As Natural Food Colorants. Thesis. Singapore: National University of Singapore.
- Linda, L. 2002. Phytosterol-esters: *Use In A Range of Food Product* [Online]. Safety and Environmental Assurance Centre. Available from: [http://www.food.gov.uk/multimedia/pdfs/d02\\_027.pdf](http://www.food.gov.uk/multimedia/pdfs/d02_027.pdf) [Accessed 17 December 2011].
- Mohammad, A., & Ahmad, S. A. 2007. Effect of Vitamin C on Blood Glucose, Serum Lipids & Serum Insulin in Type 2 Diabetes Patients. *Indian Journal of Medical Research*, **126**: 471-474.
- Mortensen, A. 2006. Carotenoids and Other Pigments as Natural Colorants. *Pure Application Chemistry*, **78** (8): 1477-1491.
- Naderi, N., Stintzing, F. C., Ghazali, H. M., Yazid, A. M., & Jazayeri, S. D. 2010. Betalain Extraction from *Hylocereus Polyrhizus* for Natural Food Coloring Purposes. *Journal of Professional Association for Cactus Development*, **12**: 143-154.
- Namugumya, B. S., & Muyanja, C. 2009. Traditional Processing, Microbiological, Physiochemical and Sensory Characteristics of *Kwete*, a Ugandan Fermented Maize Based Beverage. *African Journal of Food, Agriculture, Nutrition and Development*, **9** (4): 1046-1058.
- NSW Food Authority. 2009. *Microbiological Quality Guide for Ready-to-eat Foods* [Online]. Available from: [http://www.foodauthority.nsw.gov.au/\\_Documents/science/microbiological\\_quality\\_guide\\_for\\_RTE\\_food.pdf](http://www.foodauthority.nsw.gov.au/_Documents/science/microbiological_quality_guide_for_RTE_food.pdf) [Accessed 21 November 2011].
- Nutriweb Malaysia. 2011. *Proposed New Laws On Nutrition Labelling And Claims: What You Should Know* [Online]. Available from: <http://www.nutriweb.org.my/article.php?sid=2> [Accessed 13 December 2011].
- Ozsoy, N., Can, A., Yanardag, R., & Akev, N. 2007. Antioxidant Activity of *Smilax excels* L. Leaf Extracts. *Food Chemistry*, **110** (3): 571-583.
- Pang, K. H., Po, J. C., Chen, C. H., & Chau, C. F. 2006. Carrot Insoluble Fiber-Rich Fraction Lowers Lipid and Cholesterol Absorption in Hamsters. *Learning with Technologies: Food Science and Technology*, **30**: 337-342.
- Pascal, R. G., Glories, Y., Denis, D., & Maujean, A. 2006. *Handbook of Enology: The Chemistry of Wine Stabilization and Treatments*. Second edition. England: John Wiley and Sons, p. 59.
- Pascal, S. L., & Segal, R. 2006. Phytosterols – Biological Active Compounds in Food. *Journal of Agroalimentary Processes and Technologies*, **12** (1): 149-158.
- Patricia, B., Remedios, C., Natera, R., & Carmelo, G. B. 2006. Changes in the Polyphenolic and Volatile Content of “Fino” Sherry Wine Exposed to High Temperature and Ultraviolet and Visible Radiation. *European Food Research and Technology*, **222**: 302-309.
- Peter, C. S. 1994. *Acetic Acid* [Online]. Virginia Department of Health. Available from: <http://www.vdh.state.va.us/epidemiology/dee/publichealthtoxicology/documents/pdf/aceticacid.pdf> [Accessed 14 Decemeber 2011].
- Rebecca, O. P. S., Boyce, A. N., & Chandran, S. 2010. Pigment Identification and Antioxidant Properties of Red Dragon Fruit (*Hylocereus polyrhizus*). *African Journal of Biotechnology*, **9** (10): 1450-1454.
- Ribeiro, B, Paula, G. P., Paula, B. A., Paula, B., & Patricia, V. 2009. Fatty Acid Composition of Wild Edible Mushrooms Species: A Comparative Study. *Microchemical Journal*, **93**: 29-35.

- Rivas, A., Rodrigo, D., Martinez, A., Barbosa-Canovas, G. V., & Rodrigo, M. 2006. Effect of PEF and Heat Pasteurization on the Physical-Chemical Characteristics of Blended Orange and Carrot Juice. *LWT- Food Science and Technology*, **39** (10): 1163 – 1170.
- Roohinejad, S., Omidzadeh, A., Mirhosseini, H., Saari, N., Mustafa, S., Yusof, M., Hussin, A. S., Hamid, A., & Abdul Manap, Y. 2010. Effect of Pre-Germination Time of Brown Rice on Serum Cholesterol Levels of Hypercholesterolaemic Rats. *Journal of the Science of Food and Agriculture*, **90** (2): 245-251.
- Sampedro, F., Geveke, D. J., Fan, X., Rodrigo, D., & Zhang, Q. H. 2009. Shelf Life Study of an Orange Juice Milk Based Beverage after PEF and Thermal Processing. *Journal of Food Science*, **74** (2): 107-112.
- Saura, C. F., & Goni, L. 2009. Definition of the Mediterranean Diet Based on Bioactive Compounds. *Critical Reviews in Food Science and Nutrition*, **49** (2): 45-52.
- Shui, G., & Leong, L.P. 2002. Separation and Determination of Organic Acids and Phenolic compounds in Fruit Juices and Drinks by High-Performance Liquidchromatography. *Journal of Chromatography A*, **977**: 89-96.
- Silva, E. M., Rogez, H., & Larondelle, Y. 2007. Optimization of Extraction of Phenolics from *Ingaedulis* Leaves using Response Surface Methodology. *Separation and Purification Technology*, **55** (3): 381-387.
- Smith, J. & Edward, C. 2010. *Functional Food Product Development*. First edition. US: John Wiley and Sons, p.3.
- Somboonsilp, P., Tia, S., & Yoovidhya, T. 2011. *Development of a Small-Scale Ohmic Pasteurization System for Flavor Retention in Fruit Juice Production* [Online]. The 12<sup>th</sup> ASEAN Food Conference 2011. Available from: [http://technology.kku.ac.th/food-technology/AFC2011/index.asp\\_files/Full\\_Paper/Oral/OB\\_009.pdf](http://technology.kku.ac.th/food-technology/AFC2011/index.asp_files/Full_Paper/Oral/OB_009.pdf) [Accessed 15 December 2011].
- Soupas, 2006. *Oxidative Stability of Phytosterols in Food Models and Foods* [Online]. University of Helsinki. Available from: <http://ethesis.helsinki.fi/julkaisut/maa/skemi/vk/soupas/abstract.html> [Accessed 30 September 2010].
- Stanton, C., Ross, R. P., Fitzgerald, G. F., & Sinderen, D. V. 2005. Fermented Functional Foods Based on Probiotics and their Biogenic Metabolites. *Current Opinion in Biotechnology*, **16** (2): 198-203.
- Stoutjesdijk, P. A., Hurlestone, C., Singh, S. P., & Green, A. G. 2000. High-Oleic Acid Australian Brassica Napus and B. Juncea Varieties produced by Co-suppression of Endogenous D12-Desaturases. *Biochemical Society Transactions*, **28** (6): 938-940.
- Subramanian, G. 1995. *Quality Assurance in Environmental Monitoring: Instrumental methods*. Weinheim: VCH, p. 73-74.
- Sumbly, K. M., Grbin, P. R., & Jiranek, V. 2010. Microbial Modulation of Aromatic Esters in Wine: Current Knowledge and Future Prospects. *Food Chemistry*, **121**: 1–16.
- Taylor, C. W., & Giusti, M. M. 2009. *Optimization of the 4 Dimethylaminocinnamaldehyde Assay for Flavanols* [Online]. The Ohio State University. Available from: <https://kb.osu.edu/dspace/bitstream/handle/1811/36590/Taylor's?sequence=1> [Accessed 21 November 2011].
- Tenore, G.C., Novellino, E., & Basile, A. 2011. Nutraceutical Potential and Antioxidant Benefits of Red Pitaya (*Hylocereus polyrhizus*) Extracts. *Journal of Functional Foods*.
- Thaipong, K., Boonprakob, U., Crosby, K., Luis, C. Z., & Byrne, D. H. 2006. Comparison of ABTS, DPPH, FRAP, and ORAC Assays for Estimating Antioxidant Activity from Guave Fruit Extracts. *Journal of Food Composition and Analysis*, **19**: 660-675.
- Thoo, Y. Y., Ho, S. K., Liang, Y. L., Ho, C. W., & Tan, C. P. 2010. Effects of Binary Solvent Extraction System, Extraction Time and Extractiontemperature on Phenolic Antioxidants and Antioxidant Capacity from Mengkudu (*Morinda citrifolia*). *Food Chemistry*, **120**: 290-295.
- Treutter, D. 1989. Chemical Reaction Detection of Catechins and Proanthocyanidins with 4-dimethylaminocinnamaldehyde. *Journal of Chromatography A*, **267**: 185-193.
- Ukuku, D. O., & Sapers, G. M. 2006. Effect of Time Before Storage and Storage Temperature on Survival of *Salmonella* Inoculated on Fresh-Cut Melons. *Journal of Food Microbiology*, **24**: 288–295.

- US Food and Drug Administration. 2011. *Guidance for Industry: Juice HACCP Hazards and Controls Guidance First Edition; Final Guidance* [Online]. Available from: <http://www.fda.gov/Food/GuidanceComplianceRegulatoryInformation/GuidanceDocuments/Juice/ucm072557.htm> [Accessed 13 December 2011].
- Valentin, F., Eric, J. T., & Elizabeth, G. N. 2005. *Atherothrombosis and Coronary Artery Disease*. Second edition. US: Lippincott Williams & Wilkins, p. 40.
- Vaskonen, T., Mervaala, E., Laakso, T., & Karppanen, H. 2001. Diet Enrichment with Calcium and Magnesium Enhances the Cholesterol-Lowering Effect of Plant Sterols in Obese Zucker Rats. *Nutrition Metabolism Cardiovascular Disease*, **11** (3): 158-167.
- Vasquez, S., & Mueller, S. 2009. *Refractometer Use, Calibration and Maintenance* [Online]. University of California. Available from: <http://cemerced.ucdavis.edu/files/40574.pdf> [Accessed 21 November 2011].
- Wang, C. Y., Wu, S. C., Ng, C. C., & Shyu, Y. T. 2010. Effect of Lactobacillus- Fermented Adlay-Based Milk on Lipid Metabolism of Hamsters Fed Cholesterol-Enriched Diet. *Food Research International*, **43** (3): 819 – 824.
- Wang, M. L., Choong, Y. M., Su, N. W., & Lee, M. H. 2003. A Rapid Method for Determination of Ethanol in Alcoholic Beverages using Capillary Gas Chromatography. *Journal of Food and Drug Analysis*, **11** (2): 133-140.
- William, E. B. 2011. *Release of the Results of the AIM-HIGH Trial* [Online]. University at Buffalo Schools of Medicine and Public Health. Available from: <http://www.nhlbi.nih.gov/new/remark/boden.htm> [Accessed 22 December 2011].
- Woo, K. K., Ngou, F. H., Ngo, L. S., Soong, W. K., & Tang, P. Y. 2011. Stability of Betalain Pigment from Red Dragon Fruit (*Hylocereus polyrhizus*). *American Journal of Food Technology*, **6** (2): 140-148.
- Yang, J., Paulino, R., Janke-Stedronsky, S., & Abawi, F. 2007. Free-Radical-Scavenging Activity and Total Phenols of Noni (*Morinda citrifolia L.*) Juice and Powder in Processing and Storage. *Food Chemistry*, **102**: 302-308.
- Yang, S. C., Chen, T. I., Li, K. Y., & Tsai, T. C. 2007. Change in Phenolic Compound Content, Reductive Capacity and ACE Inhibitory Activity in Noni Juice during Traditional Fermentation. *Journal of Food and Drug Analysis*, **15** (3): 290-298.
- Yousef, A. E., & Calstrom, C. 2003. *Food Microbiology: A Laboratory Manual*. First edition. US: Wiley-IEEE, p. 5.
- Yu, L. L. 2008. *Wheat Antioxidants*. First edition. Canada: John Wiley and Sons, p. 147
- Zainoldin, K. H., & Baba, A. S. 2012. The Effect of *Hylocereus Polyrhizus* and *Hylocereus Undatus* on Physicochemical, Proteolysis, and Antioxidant Activity in Yogurt. *International Journal of Biological and Life Sciences*, **8** (2): 93-98.
- Ziaedini, A., Jafari, A., & Zakeri, A. 2010. Extraction of Antioxidants and Caffeine from Green Tea (*Camelia Sinensis*) Leaves: Kinetics and Modelling. *Food Science and Technology International*, **16**: 505-510.
- Zimmermann, B. F., & Gleichenhagen, M. 2011. The Effect of Ascorbic Acid, Citric Acid and Low pH on the Extraction of Green Tea: How to get most out of it. *Food Chemistry*, **124**: 1543-1548.