## GROWTH PERFORMANCE AND SELECTED BIOCHEMICAL INDICES ASSESSMENT IN THE SERUM AND LIVER OF *CLARIAS GARIEPINUS* JUVENILES EXPOSED TO VARYING CONCENTRATIONS OF WATER SOLUBLE FRACTION OF CRUDE OIL

## Olumide Isaac Fakolujo<sup>1</sup>, Henry Adefisayo Adewole<sup>2</sup>, Efere Martins Obuotor<sup>3</sup>, and Victor Folorunsho Olaleye<sup>2\*</sup>

<sup>1</sup>Institute of Ecology and Environmental Studies, Obafemi Awolowo University, Ile-Ife, Nigeria <sup>2</sup>Department of Zoology, Obafemi Awolowo University, Ile-Ife, Nigeria <sup>3</sup>Department of Biochemistry and Molecular Biology, Obafemi Awolowo University,Ile-Ife, Nigeria

\*Corresponding author: volaleye@oauife.edu.ng; +234-803-404-5185

**ABSTRACT.** The effect of varying concentrations (0, 0.25, 0.05 and 1.0% v/v) of water soluble fraction (WSF) of crude oil on the growth performance and biochemical changes in the serum and liver of 144 Clarias gariepinus Juveniles stocked into 12 glass aquaria at the rate of 12 fish per tank in triplicate for 90 days was assessed. The growth performance indices of the fish juvenile over the 90 days of toxic exposure revealed that the mean weight gained, specific growth rate, and total feed intake of the fish specimen decreases with increase concentration of the WSF of crude oil. However, 10% and 20% mortality were recorded in the fish exposed to 0.5% and 1.0% WSF of crude oil respectively. The activity of ALT and AST in the serum of the fish were significantly higher (p<0.05) with the increased concentration of WSF of crude oil while the ALT and AST activity of the liver of the fish juveniles irrespective of the added concentration of WSF of crude oil showed no significant differences (p>0.05). The study concluded that WSF of crude oil had adverse effects on the growth performance, blood serum and liver activities in C. gariepinus juveniles.

KEYWORDS. Crude oil, Growth performance, Serum, Enzymes, Fish, Pollution

# INTRODUCTION

The potential utility of biomarkers for monitoring both environmental quality and health of organism inhabiting polluted ecosystems has been receiving increasing attention (Gauthier *et al.*, 2004, Yusuf *et al.*, 2017).

Niger River which is one of the coastal waters in Nigeria is an important ecosystem that serves as home to 36 families and nearly 250 species of fish, of which 20 are endemic (WWF, 2006). Crude oil spillages have been reported to contribute to the decimating of rich fisheries of the Nigerian coastal waters (Adeola, 1996). Wake (2005) associated reduction in

diversity and abundance of aquatic fauna with oil-laden refinery effluents. There were also strong evidences that local fisheries are affected by migrating fish avoiding oiled areas (Powell, 1987). Ezenwa and Ayinla (1994) noted that fish recruitment for the rivers in the Delta Area of Nigeria has been drastically reduced and this factor accounts for the poor catch of fish by the artisanal and pelagic fishermen, thereby leading to low food supply and income.

Generally, crude oil has been reported to be primarily toxic to aquatic organisms due to the presence of Polycyclic Aromatic Hydrocarbons (PAHs) (Heintz *et al.*, 1999). Baden, (1982) reported significantly increase in oxygen consume soluble fraction (WSF). Mortality and oxidative stress were observed in tilapia fingerlings exposed to the WSF of diesel fuel (Dede and Kaglo, 2001) and growth and feed utilization were significantly reduced in exposed *Heterobrachus bidorsalis* (Ofojekwu and Onah, 2002). Drastic changes in liver enzyme activities of the catfish, *Clarias gariepinus* were also reported following exposure to crude oil (Sunmonu and Oloyede, 2006).

However, the viscosity of oil in aquatic environment coupled with the presence of hydrocarbons in oil poses a carcinogenic threat to fish, birds and mammals (Mason, 1996). Oil spillage over time can affect species populations through the destruction of more sensitive juvenile stages (Horsefall and Spiff, 1998) and is capable of sublethal stress effects, mutation and changes in behavioural patterns in individual organisms.

Aquatic environments are made up of complex interactions between plants and animal species and their physical environment. Harm to the physical environment will often lead to physiological responses (such as unpleasant smell and taste of the toxicant, lower food intake, incessant jumping, gulping of air, restlessness, loss of equilibrium, increased opercular activities, surface to bottom movement, sudden quick movement, and resting at the bottom) which can result to death of one or more species in a food chain leading to damage for other species further up the chain (Sunmonu, 2006; Ezemoye and Ogbomida, 2010).

Fish which is a cheap source of animal protein play important role in the human diet. Therefore, the vulnerability of the *Clarias gariepinus* juvenile to the toxic effects of water soluble fractions of crude oil and its refined products in order to prevent the decimating rich fisheries in the Nigerian Coastal waters needs to be evaluated, hence this study. Growth Performance and Selected Biochemical Indices Assessment in The Serum and Liver of Clarias Gariepinus Juveniles Exposed to Varying Concentrations of Water Soluble Fraction of Crude Oil

# **MATERIALS & METHODS**

## Water Soluble Fraction (WSF) Preparation

Fifteen (15) litres of Excravos-grade crude oil were sourced from the Department of Petroleum Resources, Warri, Delta State, Nigeria. Volumes of WSF of crude oil used for this experiment were prepared according to the method of Afolabi *et al.* (1985) under dim illumination. 200 ml of Crude oil was added to 800 ml of distilled water in a capacity borosilicate screw-capped conical flask. The mixture was shaken for 24 hrs using a Gallenkamp orbital stirrer and was allowed to stand for at least 3 hours after which it was poured into a glass-stoppered separating funnel. The mixture was further allowed to settle over night to allow the oil droplets in the mixture to settle in the upper layer leaving the clear WSF in the lower part of the separating funnel. The pure WSF was then siphoned into a dark coloured screw-capped Winchester bottle until needed for the experiment.

## Fish Stocking and Exposure to WSF of Crude oil

Two hundred (200) six-week old juvenile *Clarias gariepinus* which were obtained from BODMET fish farm, Ile-Ife, Osun State, Nigeria were acclimatized to laboratory conditions in Fish Culture laboratory, Department of Zoology, Obafemi Awolowo University, Ile-Ife, Nigeria for 14-days and fed *ad libitum* with 2 mm Durantee floating feed. After acclimatization, 144 fish with a mean weight of  $43.34 \pm 0.78$  g were randomly selected and stocked in 12 glass aquaria (30 cm x 60 cm x 45 cm), each containing 30 litres of water at the rate of 12 fish per tank. WSF at 0% (control), 0.25%, 0.5%, and 1% concentrations were introduced into each tank in triplicates. The fish were fed at 3% of the body weight twice daily (Adewolu and Adeoti, 2010) for 90 days.

## **Data Collection**

The initial weight of the fish were taken before the commencement of the experiment and biweekly subsequently. On-biweekly weighing days, crushed ice was added into water to reduce the fish activity before being removed for weighing after a couple of minutes. The ration allotted to the fish was adjusted based on the new weight gain by the fish after each weighing. The aggregate weight of the test fish per tank was used to calculate the fish growth performance indices as described by Sveier *et al.* (2000):

Mean weight gain (MWG) =  $(W_f-W_i)$  n; Where;  $W_i$  = initial body weight  $W_f$  = final body weight n=number of fish. Specific Growth rate (SGR) =  $(Log W_f-Log W_i)$  t x 100 Food Conversion ratio (FCR) = Feed Intake(g) / Weight gain Total Food Intake (TFI) = Total weight of the fish/ Total number of fish in the tank Survival rate= Number of fish at the end of experiment / number of fish stocked x 100

### Serum and Liver Biochemical Assays

After 90 days of the experiment, blood sample for serum analysis was obtained through the caudal puncture from 5 randomly selected fish from each treatment using a disposable hypodermic syringe and needle. The blood sample was expressed into plain specimen bottle and was allowed to clot, then centrifuge at 3500 rpm for 15 mins to separate blood cells from the serum. This was then collected into another plain bottle and stored at -4°C in a Thermocool deep freezer for the evaluation of Aspartate transaminase and alanine transaminase. The blood serum of the fish was analyzed for the activity of AST and ALT spectrophotometrically according to Reitman and Frankel (1957) and Schmidt and Schmidt (1963).

One (1) gram of the excised liver of the test fish was homogenized in a 10 ml Tris buffer solution using rotor of a Tri-R Stir-R electronic homogenizer. The homogenate samples were then assayed using spectrophotometer for the levels of Aspartate Transaminase (AST), Alanine Transaminase (ALT) and Lactate Dehydrogenase (LDH) following the methods of Reitman and Frankel (1957), Schmidt and Schmidt (1963) and Weisshaar et al. (1995) respectively.

#### RESULTS

#### **Growth Performance and Feed Utilization**

The bi-weekly growth performance of the *C. gariepinus* juveniles exposed to varying concentrations of WSF of crude oil is shown in Figure 1. The fish in the control treatment had the better growth performance. The growth of the fish juvenile exposed to varying concentration of WSF of crude oil was observed to be retarding at the sixth week of the experiment, although a sudden rise was observed in the growth of the fish specimens exposed to 0.05% WSF of crude oil at the tenth week, but was however later retarded in the subsequent weeks. The growth performance indices of the fish juvenile over 90 days of toxic exposure, as shown in Table 1, revealed that the mean weight gained, specific growth rate, and total feed intake of the fish specimens which ranged between 4.80  $\pm$ 1.06 g (1.00% WSF) to 62.62  $\pm$ 2.25 g (0% WSF (Control)); 0.05  $\pm$ 2.25% (1.00% WSF) to 0.45  $\pm$ 2.90 (0% WSF), and 13.77 (1.00% WSF) to 16.37 (0% WSF) respectively were statistically different (p<0.05) and was also observed to decrease with increased concentration of the WSF of crude oil.

Growth Performance and Selected Biochemical Indices Assessment in The Serum and Liver of Clarias Gariepinus Juveniles Exposed to Varying Concentrations of Water Soluble Fraction of Crude Oil



Figure 1: The bi-weekly growth performance of the *C. gariepinus* juveniles exposed to varying concentrations of WSF of crude oil.

The FCR of the fish exposed to WSF of crude oil were 3 or more folds than the FCR of the fish in control treatment.

No mortality was recorded in the control (0% WSF) culture tank and in the fish exposed to 0.25% WSF of crude oil. However, 10% and 20% mortality were recorded when the fish juveniles were exposed to 0.50% and 1.00% WSF of crude oil respectively (Table 1).

## **Enzyme Activities**

The summary of assayed hepatic enzyme activities in the blood serum of *C. gariepinus* juveniles exposed to WSF of crude oil and the control is shown in Table 2. ALT activity in the serum of the fish juveniles exposed to 1.00% WSF ( $30.98 \pm 1.52$  U/L) was significantly higher (p<0.05) than in the fish juveniles within the control culture tank and those exposed to lower concentrations of crude oil. Generally, the trend showed that the higher the concentration of the WSF of crude oil, the higher the serum ALT concentration. Analysis also showed that exposure of the fish juveniles to WSF of crude oil significantly (p<0.05) increased the hepatic ALT concentration (Table 2).

Growth	Concentration of WSF				
Performance Indices	0% (Control)	0.25%	0.50%	1.00%	
Initial mean	40.93 <sup>a</sup>	42.65 <sup>a</sup>	43.41 <sup>a</sup>	46.37 <sup>a</sup>	
weight (g)	$\pm 0.52$	$\pm 0.32$	±0.63	$\pm 0.92$	
Final mean	103.53 <sup>a</sup>	61.26 <sup>b</sup>	59.81 <sup>b</sup>	51.17 <sup>b</sup>	
weight (g)	$\pm 5.28$	±3.12	$\pm 2.26$	$\pm 3.58$	
Mean weight	62.61 <sup>a</sup>	18.61 <sup>b</sup>	16.40 <sup>b</sup>	4.80 <sup>c</sup>	
gain (g)	$\pm 2.25$	$\pm 2.44$	±3.49	$\pm 1.06$	
Mean Daily	$0.70^{a}$	0.21 <sup>b</sup>	0.18 <sup>b</sup>	0.05 <sup>c</sup>	
Weight Gain (g)	±0.30	$\pm 0.17$	±0.15	$\pm 0.40$	
Specific Growth	$0.45^{a}$	0.18 <sup>b</sup>	0.15 <sup>b</sup>	0.05 <sup>c</sup>	
<b>Rate</b> (%)	$\pm 0.96$	$\pm 0.72$	$\pm 0.45$	±0.25	
Total Feed Intake (TFI) (g)	16.37	14.16	13.86	13.77	
Feed Conversion	$0.26^{a}$	0.76 <sup>b</sup>	0.85 <sup>b</sup>	2.87 <sup>c</sup>	
Ratio (FCR)	$\pm 0.08$	±0.37	$\pm 0.60$	±1.41	
% Mortality	0	0	10	20	

# **Table 1:** Mean Growth Performance\* of C. gariepinus Juveniles Exposed to Different<br/>Concentrations of WSF of Crude Oil for 90 days

\*Values with different superscripts are significantly different at p<0.05

# **Table 2:** Serum\* ALT and AST Levels in *C. gariepinus* Juveniles Exposed to Different Concentrations of WSF of Crude Oil

WSF	Enzyme Activity (U/L)			
Concentration	Alanine Transaminase (ALT)	Aspartate Transaminase (AST)		
0%	$9.60^{\circ} \pm 0.30$	448.00 <sup>c</sup> ±14.07		
0.25%	$9.67^{c} \pm 1.02$	497.45 <sup>bc</sup> ±11.74		
0.50%	17.51 <sup>b</sup> ±0.57 537.09 <sup>b</sup> ±20.16			
1.00%	$30.98^{a}\pm 1.52$	$612.20^{a}\pm 18.77$		

\*Values with different superscripts are significantly different at p<0.05

Exposure of *C. gariepinus* juveniles to WSF of crude oil significantly (p<0.05) increased the activity of AST (Table 2). The results showed that AST activity in the serum of fish juveniles exposed to 1% WSF of crude oil ( $612.20 \pm 18.77 \text{ U/L}$ ) was significantly higher (p<0.05) than in the control and test fish exposed to 0.25% and 0.50% of WSF of crude oil.

As shown in Table 3, the ALT activity in the liver of fish juveniles irrespective of the added concentration of WSF of crude oil showed no significant difference (p>0.05). Nonetheless, ALT activity in the liver of fish juveniles exposed to 1.00% WSF (70.50  $\pm$ 1.40 U/L) was correspondingly lower than in 0.50% WSF (81.25  $\pm$ 5.80 U/L).

Similarly, AST activity in the liver was lowest in the fish juvenile exposed to 1.00% WSF of crude oil (399.13 ±17.18 U/L) (Table 3) and the enzyme activity was also observed to decrease in the liver of the fish with higher concentrations of WSF of crude oil exposure, though not significantly.

Hepatic lactate dehydrogenase (LDH) activity was lowest (294.05  $\pm$ 72.48 U/L) in fish juveniles exposed to 0.25% WSF of crude oil was highest (705.72  $\pm$ 28.54 U/L) in the fish exposed to 0.05% WSF. Analyses however showed that the activity of the enzyme in the fish exposed to 1.00% WSF of crude oil was not significantly different (p<0.05) from those of the fish in the control tank (Table 3).

**Table 3:** ALT, AST and LDH Levels\* in the Liver of C. gariepinus Juveniles Exposed toDifferent Concentrations of WSF of Crude Oil

Enzyme Activity (U/L)						
WSF Concentration	Alanine Transaminase (ALT)	Aspartate Transaminase (AST)	Lactate Dehydrogenase (LDH)			
0%	$42.75^{b} \pm 6.83$	439.33 <sup>a</sup> ±35.57	478.73 <sup>ab</sup> ±87.58			
0.25%	$79.96^{a}\pm 6.96$	$476.40^{a} \pm 31.46$	$294.05^{b}\pm72.48$			
0.50%	$81.25^{a} \pm 5.80$	$432.29^{a} \pm 14.94$	705.72 <sup>a</sup> ±28.54			
1.00%	$70.50^{a} \pm 1.40$	399.13 <sup>a</sup> ±17.18	$538.16^{ab} \pm 42.68$			

\*Values with different superscripts are significantly different at p<0.05

## DISCUSSION

Better growth performances were observed in the *C. gariepinus* juveniles within the control (0% WSF) compared to the test fishes exposed to varying concentrations of WSF of crude oil in this study. Poor growth performances in the exposed group could be attributed to the effects of the soluble fractions of crude Polycyclic Aromatic Hydrocarbon (PAH) and Total Petroleum Hydrocarbon (TPH) which probably affected the fish eco-physiology (Hodson *et al.*, 1997). The high feed conversion ratio (FCR) observed in the exposed group indicated the inability of the exposed fish to convert feed consumed into required body protein (Sunmonu and Oloyede, 2006). Low mean weight gain recorded in the exposed fish might also be due to the immediate use up of protein content in feeds through oxidative stress rather than being utilized for new protein formation (Murty *et al.*, 1983).

However, relatively high mortality through possible disease contagion observed in test fish exposed to 0.5% and 1.0% WSF concentration of crude oil was due to the potency of environmental toxicants which increased disease susceptibility by interfering with immune, reproductive and developmental processes within aquatic animals (Couch and John, 1978). Excess mucus secretion observed in test fish was probably a neurological response to the irritation caused by test fish exposure to WSF of crude oil. Wang (1986) also reported that excessive mucus secretion could lead to the thickening of test fish skin or its gill epithelia which might result in fish mortality.

Growth Performance and Selected Biochemical Indices Assessment in The Serum and Liver of Clarias Gariepinus Juveniles Exposed to Varying Concentrations of Water Soluble Fraction of Crude Oil

Exposure to varying concentrations of WSF of crude oil was found to induce some biochemical changes in the liver and blood enzymes in *C. gariepinus* juveniles. Alanine transaminase (ALT) activity was found to increase in the serum of the exposed test fish with the enzyme activity level increasing with succeeding concentrations of WSF of crude oil. Aspartate transaminase (AST) activity level in the serum of the exposed test fish especially those exposed to 0.5% and 1.0% WSF of crude oil were significantly higher (p<0.05) than those of the control fish (0% WSF). These observations corroborated the results of Lin *et al.*, 2002); Palanivelu *et al.*, 2005); Abdel-Moneim *et al.*, (2008) and Wegwu and Omeodu (2010). According to the authors, hepatotoxicity of WSF of crude oil probably caused cell membrane damage where hepatic enzymes are emptied into the blood stream which probably resulted in the elevation of ALT/AST activity in the serum. However, the decreasing hepatic AST activity with increasing concentration of WSF of crude oil was attributed to biochemical changes due to the environmental stress resultant from exposure to the toxicants (Abdel-Moneim *et al.*, 2008)

Val and Almeid –Val (1999) reported that LDH activity increase in the liver was able to catalyse the conversion of pyruvate to lactate anaerobically. LDH activity recorded in this study was highest in the liver of *C. gariepinus* juveniles exposed to 0.5% and 1.0% WSF of crude oil. This may be due to low dissolved oxygen uptake which occurred in test fish under toxic exposure (Sunmonu and Oloyede, 2006) and could also be due to the presence of liver leisons which might have compromised its structural integrity (Ramesh *et al.*, 1993; Das *et al.*, 2004 a,b)

## CONCLUSION

The study concluded that WSF of crude oil had adverse effects on the growth performance, blood serum and liver activities in *Clarias gariepinus* juveniles.

### REFERENCES

- Abdel-Moneim AM, Abou Shabana NM, Khadre SEM, Abdel-Kader HH. Physiological and histological effects in catfish (*Clarias lazera*) exposed to dyestuff and chemical wastewater. International Journal of Zoological Research 2008; 4:189-202
- Adeola MO. Present and potential Sources of Water Pollution Effects on Fish and Wildlife Resources. In Aina E.O.A. and Adefioye, N.O. (Eds). FEPA Monographs 1996; 6:191-197
- Adewolu MA, Adeoti AJ. Effect of mixed feeding schedules with varying dietary crude protein levels on the growth and feeding utilization of *Clarias gariepinus* (Burchell, 1822) fingerlings. Journal of Fisheries and Aquatic Science 2010; 5:304-310
- Afolabi OA, Adeyemi SA, Imevbore AMA. Studies on toxicity of some toxicity of some Nigeria crude oils to some aquatic organisms. In: Proceedings of 1985 Int. Seminar on the Petroleum Industry and the Nigeria Environment. NNPC, Lagos: 1985; 269-273
- Baden SP. Oxygen consumption rate of shrimp exposed to crude oil extract. Marine Pollution Bulletin 1982; 13(7):230-233
- Couch C, John A. Disease, Parasites, and toxic responses of commercial penaeid shrimps of the Gulf of Mexico and South Atlantic coast of North America. EPA-600/J-78-072. U.S. Fish and Wildlife Service. Fishery Bulletin 1978; 76:1-44.
- Das P, Ayyapan S, Das B, Jena J. Nitrite toxicity in Indian Major carps: sublethal effect on selected enzymes in fingerlings of *Catla catla, Labeo rohita* and *Cirrhinus mrigala*. Comparative Biochemistry and Physiology C-Toxicology and Pharmacology 2004a; 138:3-10
- Das P, Ayyapan S, Jena J, Das B.. Acute toxicity of ammonia and its sub-lethal effects on selected heamatological and enzymatic parameters of mrigal, *Cirrhinus mrigala* (Hamilton). Aquaculture Research 2004b; 35:134-143
- Dede EB, Kaglo HO. Aqua-toxicological effects of water soluble fraction (WSF) of diesel fuel on *Oreochromis niloticus* fingerlings. Journal of Applied Sciences and Environmental Management 2001; 5(1):93-96
- Ezenmoye L, Ogbomida TE. Histopathological effects of gammalin 20 on African catfish (*Clarias gariepinus*) Applied and Environmental Soil Science (2010); 1-8
- Ezenwa BI, Ayinla OA. Conservation strategies for endangered fish breeding and nursery grounds within the coastal wetlands of Nigeria. Aquatic Conservation, Marine and Fresh Water Ecosystems 1994; 4:125-133
- Gauthier L, Tardy E, Mauchet F, Marty J. Biomonitoring of the genotoxic potential (micronucleus assay) and detoxifying activity (EROD induction) in the River Dadau (France), using the amphibian *Xenopus laevis*. Scientific Total Environment 2004; 323:47-61
- Heintz RA, Short JW, Rice SD. Sensitivity of fish embryos to weathered crude oil Part II. Increased mortality of pink salmon (*Onchorpynchus gorbuscha*) Environmental Toxicology and Chemistry 1999; 18(3):493-503

- Hodson PV, Maj MK, Efler S, Burnison BK, Van Heiningen ARP, Girard R, Carey JH. Experiment with oil and dispersants on the sea grass *Zostera holtii*. Report of the Advisory Committee on Pollution of the Sea, Field Study Council London. 1997
- Horsefall M, Spiff AI. *Principles of Environmental Chemistry*, Metropolis Ltd., Port Harcourt .1998; p 112-119.
- Lin SC, Chung TC, Ueng TH, Lin YH, Hsu SH, Chiang CL, Lin CC. The hepatoprotective effects of *Solanum alatum* Moench on acetaminophen-induced hepatoxicity in mice. A.M.J. Uin. Med. 2002; 28:105-114
- Mason CF. Biology of Freswater Pollution, 3<sup>rd</sup> Ed., Longman and Scientific Technical, Harlow, U.K. 1996; 356 pp
- Murty AS, Rajabhushanam BR, Ramani AV, Christopher K. Toxicity of fenitrothion to the fish, *Mystus cavasius* and *Labeo rohita*. Environmental Pollution Series A: 1983; 30:225-232.
- Ofojekwu PC, Onah JA. The effects of water soluble fraction of crude oil on the growth of the catfish, *Clarias gariepinus* (Burchell, 1822). African Journal of Environmental Pollution and Health 2002; 1(2):1-7
- Palanivelu V, Vijayavel K, Ezhilarasibalasubramanian S, Balasubramanian MP. Influence of insecticidal derivative (Cartap Hydrochloride) from the marine polychaete on certain enzyme systems of the freshwater fish *Oreochromis mossambicus*. Journal of Environmental Biology 2005; 26:191-196
- Powell CB. Effects of freshwater oil spillages on fish and fisheries. *In: proceedings of 1987* Seminar on the petroleum Industry and the Nigerian Environment. NNPC, Lagos 1987; 208-227
- Ramesh M, Sivakumari K, Kanagaraj M. Toxicity of dye effluent in lactate dehydrogenase activity in *Labeo rohita. Journal of Environmental Protection* 1993; 13:124-127
- Reitman S, Frankel S. A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminase. American Journal of Clinical Pathology 1957; 28(1):56-63
- Schmidt E, Schmidt FW. Determination of serum GOT and GPT. Enzym. Biol. Clin. 1963;
   3:1Sunmonnu TO. Biochemical effect of crude oil polluted water on catfish (*Clarias gariepinus*) and the effect of its consumption on selected rat cellular system. Ph.D. thesis, Department of Biochemistry, University of Ilorin, Ilorin: 2006; 58pp
- Sunmonu TO, Oloyede OB. Changes in liver enzyme activities in African catfish (*Clarias gariepinus*) Exposed to Crude Oil Exposure. Asian Fish Science 2006; 19:104-112.
- Sveier H, Raae AJ, Lied E. Growth and protein turnover in Atlantic salmon (*Salmo salar*): The effect of dietary protein level and protein particle size. Aquaculture 2000; 185:101-120.
- Val AL, Almeid-Val VMF. Effects of crude oil on respiratory aspects of some fish species of the Amazon. In: *Biology of Tropical Fish (Ed. A.L. Val and V.M.F. Almeida-Val)*. National Institute for Research in the Amazon, Manaus: 1999; 227-291

- Wake H. "Oil Refineries: A review of their ecological impacts on the aquatic environment". Estuarine, Coastal and Shelf Science 2005; 62:131-140
- Wang W. Toxicity test of aquatic pollutants by using common duckweed environmental pollutant. Environ. Poll. 1986; 11:1-14.
- Wegwu MO, Omeodu SI. Evaluation of selected biochemical indices in *Clarias gariepinus* exposed to aqueous extract of Nigerian crude oil (Bonny Light). Journal of Applied Science and Environment Management. 2010; 14(1):77-81
- Weisshaar D, Gossrau E, Faderl B. Normal ranges of alpha-HBDH, LDH, AP and LAP as measured with substrate optimated test charges. Med Welt 1975; 26:387.
- WWF (World Wildlife Fund) Fishing on the Niger River. 2006; Retrieved May 10, 2007 from

http://www.panda.org/news\_facts/multimedia/video/index.cfm?uNewsID=61121.

Yusuf OA, Adewole HA, Olaleye VF. Assessment of the Water Quality of Saba River, Osogbo, Nigeria. Notulae Scientia Biologicae 2017; 9(2):188-195.